

A New Perturbation Theory of Finite-Size Effects near Critical Point

C.B. Yang^{1,2} and X. Cai¹

¹*CCAST(World Laboratory), P.O. Box 8730, Beijing 100080, China*

²*Institute of Particle Physics, Hua-Zhong Normal University, Wuhan 430070, China**

()

Abstract

A new perturbation theory is proposed for studying finite-size effects near critical point of the ϕ^4 model with a one-component order parameter. The new approach is based on the techniques of generating functional and functional derivative with respect to external source field and can be used for temperatures both above and below the critical point of the bulk system. It is shown that this approach is much simpler comparing with available perturbation theories. Particularly, this new method avoids renormalization in calculating many physical quantities such as correlation functions etc..

PACS numbers: 68.35.Rh, 75.10.Hk, 05.70.Jk

Typeset using REVTeX

*Mailing address

Finite-size effects near critical points have been remained over the past two decades to be an important topic of the active research both theoretically and experimentally [1]. When one or more dimensions of a bulk system is reduced to near or below a certain characteristic length scale, the associated properties are modified reflecting the lower dimensionality. It is believed that finite-size effects are precursors of the critical behavior of the infinite system and can be exploited to extract the limiting behavior. A central role plays the finite-size scaling behavior predicted by both the phenomenological [2] and renormalization group [3] theories. Those theories allowed a systematic discussion of the finite-size effects and, consequently, form the cornerstone of our current understanding of the way in which the singularities of an infinite system are modified by the finiteness of the system in some or all of the dimensions. Of course, the exact form of scaling functions can't be given in those scaling theories.

In 1985, Brézin and Zinn-Justin (BZ) [4] and Rudnick, Guo and Jasnow (RGJ) [5] developed two field-theoretical perturbation theories for the calculation of the finite-size scaling functions within the ϕ^4 model which corresponds to the Ising model. Most applications of these theories to three-dimensional systems have been restricted to region $T \geq T_C$ [6] with a few calculations in region below T_C [7]. In recent years the ϕ^4 and the extended ϕ^6 models have been used to investigate the multiplicity fluctuations in the final states for first- and second-order phase transitions of quark gluon plasma [8], under the approximation similar to the so-called zero-mode approximation. As pointed out in the first paper in Ref. [9], the theory of BZ is not applicable for $T < T_C$ and the results from RGJ theory are not quantitatively reliable in the same temperature region since the coefficients of the Gaussian terms in the integrals are negative for those temperatures. Although the modified perturbation method in Ref. [9] can be used for both $T > T_C$ and $T < T_C$, the calculation is lengthy and can be done only to the first-order in practice. Since one does not know the exact order of values of higher order terms, theoretical results have large uncertainty. Furthermore, in the calculations of Ref. [4-7,9] there are divergent terms which must be treated using renormalization technique which not only is complicated but also increases theoretical uncertainty.

Therefore, a further study of the perturbation theory of finite-size effect is necessary.

It should be pointed out that all perturbation theories mentioned above are based on the Fourier decomposition of the order parameter. This method is natural because the decomposition enables one to transform the functional integral into an infinite product of tractable normal integrals. Although such decomposition has simple physical explanation which is very fruitful for the understanding of properties of infinite systems and can deduce reliable physical results, as in the case of usual field theories in particle physics, it brings about a great deal of calculations for systems with finite-size. This is not surprising. As is well-known, quantities complicated in coordinate space may have simple momentum spectra thus looks simple in momentum space, but those obviously nonzero only in a finite range must have puzzling momentum spectra. Therefore, for the study of properties of finite-size systems, calculations in coordinate space might be simpler and more effective. The point here is that one must calculate the complicated functional integral which is very difficult to be evaluated directly.

In this Letter, we employ the technique of generating functional with the help of external source field to develop a perturbation theory for the study of the finite-size effects of systems near their critical points. The external source field enables one to turn the functional integral into much easier functional derivatives, thus facilitates the calculations greatly. As will be shown, the perturbation expansion can be carried out easily and one can get expressions for physical quantities to second-order with little exertion. More importantly, such technique isolates all the divergence in every order of the expansion for the partition function into a common factor, showing the common source of those divergences in the calculating of thermodynamical quantities. Because of this factorization the renormalization can be done only once for the perturbative calculations so that the expansion can be done to high order needed with special difficulties. This advantage will make more detailed investigation of finite-size effects possible.

In a ϕ^4 model of phase transition with a one-component order parameter, the partition function which is most important for the calculation of other thermodynamical quantities

can be expressed as a functional integral of exponential of the Hamiltonian H of the system

$$Z = \int \mathcal{D}\phi \exp(-H) = \int \mathcal{D}\phi \exp \left\{ - \int d^d x \left[\frac{\gamma}{2} \phi^2 + \frac{1}{2} (\nabla \phi)^2 + \frac{u}{4!} \phi^4 \right] \right\} , \quad (1)$$

in which $\gamma = a(T - T_C)$, a and u are temperature dependent positive constants, T_C is the bulk critical temperature, ϕ is the order parameter of the system, and the integration over x is in the range of the finite volume.

Let's first consider the calculation of partition function in the region $T > T_C$ i.e. $\gamma > 0$. To calculate the functional integral in Eq. (1), one can begin with considering a Gaussian functional integral with the introduction of external source field J

$$\begin{aligned} W[J] &= \int \mathcal{D}\phi \exp \left\{ - \int d^d x \left[\frac{\gamma}{2} \phi^2 + \frac{1}{2} (\nabla \phi)^2 - J\phi \right] \right\} \\ &= \int \mathcal{D}\phi \exp \left\{ - \int d^d x \left[\frac{\gamma}{2} \phi^2 - \frac{1}{2} \phi \nabla^2 \phi - J\phi \right] \right\} , \end{aligned} \quad (2)$$

where it is assumed that the $(d-1)$ -dimensional surface integral of $\phi \nabla \phi$ equals to zero which is satisfied for the usually assumed periodic, anti-periodic, and Dirichlet boundary conditions. Following the same standard procedures as in field theory for infinite system, one has

$$W[J] = [\det(\gamma - \nabla^2)]^{-1/2} \exp \left[\frac{1}{2} \int d^d x d^d y J(x) \Delta(x, y) J(y) \right] , \quad (3)$$

where $\det$ represents the determinant of the operator $\gamma - \nabla^2$ and the Green's function $\Delta(x, y)$ is the inverse of the same operator, namely $\Delta(x, y)$ satisfies

$$(\gamma - \nabla^2) \Delta(x, y) = \delta(x - y) . \quad (4)$$

The boundary condition for $\Delta(x, y)$ can be chosen to be Dirichlet type. This Green's function can be obtained analytically for some systems with regular boundaries. For the case of the simplest one-dimensional system within $(0, L)$, one can show with a few algorithms that $\Delta(x, y)$ is quite simple and can be written as

$$\Delta(x, y) = \begin{cases} \sinh \omega(L - y) \sinh \omega x / (\omega \sinh \omega L) & \text{for } x < y, \\ \sinh \omega(L - x) \sinh \omega y / (\omega \sinh \omega L) & \text{for } x > y, \end{cases} \quad (5)$$

in which $\omega = \sqrt{\gamma}$, $\sinh x \equiv \frac{1}{2}(\exp(x) - \exp(-x))$ is the hyperbolic sine function.

Different from those for infinite systems, this Green's function for finite-size system can't be written as a function of single variable $x - y$ even in the case with periodic boundary condition. The usually assumed translational invariance is destroyed generally due to of the finite size of the system and the system is invariant only under a subgroup of the translational transformation under the assumption of periodic boundary condition. It should also be pointed out that the last expression for the Green's function can be continued to the temperature $T = T_C$, so that the Green's function for $\gamma = 0$ will be treated as the limiting of $\gamma \rightarrow 0_+$ and will not be discussed in this Letter.

In terms of $W[J]$, the partition function Z can be expressed as

$$\begin{aligned} Z &= \exp\left(-\frac{u}{4!} \int d^d x \frac{\delta^4}{\delta J^4(x)}\right) W[J] \Big|_{J=0} \\ &= \sum_{n=0}^{\infty} \frac{(-u)^n}{n!(4!)^n} \left[\int d^d x \frac{\delta^4}{\delta J^4(x)} \right]^n W[J] \Big|_{J=0} \end{aligned} \quad (6)$$

Owing to the fact that it involves only functional derivatives, the last expansion can be evaluated quite easily with the Wick's theorem. In traditional perturbation theories, Z can only be obtained up to the first-order. In our new approach, the partition function can be calculated to higher order and, for example, takes the form up to the second-order approximation,

$$\begin{aligned} Z &= [\det(\gamma - \nabla^2)]^{-1/2} \left[1 - \frac{u}{8} \int d^d x \Delta^2(x, x) + \frac{u^2}{8} \int d^d x d^d y \right. \\ &\quad \left. \left(\frac{\Delta^2(x, x)\Delta^2(y, y)}{8} + \Delta(x, x)\Delta^2(x, y)\Delta(y, y) + \frac{\Delta^4(x, y)}{3} \right) \right]. \end{aligned} \quad (7)$$

Because no singularity is associated with the Green's function $\Delta(x, y)$, a normal integral will give the result. In some cases for which the Green's function has a simple form, above integral can be carried out analytically. It should be noted that all terms in the perturbation expansion are finite except a common factor $[\det(\gamma - \nabla^2)]^{-1/2}$. This factor depends on the boundary condition and may contribute to thermodynamical quantities such as entropy and heat capacity etc.. The factor is generally divergent, therefore renormalization is needed to

obtain physically acceptable results. The renormalization for the factor can be done with ϵ -expansion and/or in fixed dimensions, as shown in Ref. [4-7,9]. So we do not discuss the renormalization of the factor in this Letter. Since all the divergences have been isolated into the common factor, the renormalization needs to be done only once, so that one can expand the series to any order he needs with little effort. In fact, to every next expansion order, the terms are accompanied with integration of two more Green's functions over the volume of the system. As can be seen from an example of the expression for the Green's function in one-dimensional case, The integral of the two more Green's function is quite small, thus the first a few terms in the expansion is enough in most cases, though higher order calculations are not very difficult. More interestingly, the common divergent factor is cancelled and needs not to be worried about when one is interested in non-thermodynamical quantities such as correlation functions and moments of the order parameter, leaving those physical quantities finite at any order of the perturbation expansion. It is clearly seen from the symmetry property of the Hamiltonian that correlation functions involving odd number points are always zero so that only correlation functions involving even number points are nonzero and finite. For illustration, the two-point correlation function is calculated up to first-order approximation

$$\langle \phi(x)\phi(y) \rangle = \Delta(x, y) - \frac{u}{4} \int d^d z \Delta(x, z) \Delta(z, z) \Delta(z, y) . \quad (8)$$

One can see that no singularity exists in last expression and its extension to higher orders is straightforward. The same is true for other non-thermodynamical quantities such as moments of the order parameter etc.. Due to the violation of translational invariance, the two-point correlation function depends the coordinates x, y of the two points separately instead on their difference $x - y$. That is to say, $\langle \phi(x)\phi(y) \rangle \neq \langle \phi(x - y)\phi(0) \rangle$, contrary to the usual assumption of translational invariance, except in the very central region of the system. This is different from traditional perturbation theories for finite-size systems.

From above discussions one sees that the forms of the expansion are the same for systems in one-, two- and three-dimensional spaces. To deal with spaces with different dimensions,

one needs only to use a new Green's function in proper dimensional space and retains the form of the expansion. More remarkably, if the shape of the boundary of the system is not a square nor cubic, the Fourier decomposition can't be performed and the traditional perturbation theories fail to work. This disadvantage limits the application of traditional perturbation theories to realistic systems for which the boundaries are usually not so regular. For these systems our new approach can nevertheless be carried out because the needed Green's function and the determinant of the operator $\gamma - \nabla^2$ can still be determined numerically while the form of our perturbation expansion needs not to be altered. This advantage will offer the possibility for one to study more realistic models. It is certain that the comparison between results from the new perturbation theory and real experiments can be done in the future.

When one intends to calculate quantities in the temperature region lower than T_C , above procedures should be modified. In that temperature region γ is negative so that the functional integral in Eq. (2) is ill-defined and can't be used in the calculations of the partition function Z and other physical quantities. In fact, this is the very origin of all the difficulties in previous perturbation theories. The cause for the difficulties can be understood physically. For $T > T_C$, the potential $(\gamma/2)\phi^2 + (u/4!)\phi^4$ takes its minimum at $\phi = 0$ and fluctuations about the minimum should be small and can be taken into account by perturbation theory with the help of $W[J]$ as shown above. For $T < T_C$, however, the potential takes its extremum at $\phi = 0$ and its degenerate minima at $\phi = \pm\phi_0 = \pm\sqrt{6a(T_C - T)/u}$. These extrema can be easily understood by drawing the potential for this case. That is to say, the most probable order parameter or ground state of the system for temperature below T_C is either ϕ_0 or $-\phi_0$. The ground states do not possess the symmetry $\phi \leftrightarrow -\phi$ of the Lagrangian, or the symmetry is broken. This is known as the spontaneous symmetry breaking which is very familiar in superconductors, crystal lattices, and the buckling of a compressed needle. In such case, $\langle\phi\rangle$ is nonzero but lies in the vicinity of either ϕ_0 or $-\phi_0$. If the perturbation expansion is nevertheless be done around $\phi = 0$, it is bound to be very complicated and even to fail. Then we are authorized to expand perturbation theory around

one of the degenerate classical minima of the potential. Keeping all above facts in mind, one can shift the field to the vicinity of one of the minima through, for example,

$$\phi = \phi_0 + \varphi \quad (9)$$

and considers fluctuations of φ around zero instead of those of ϕ . Then the Hamiltonian of the system takes the form

$$H = \int d^d x \left[-\frac{3\gamma^2}{2u} + \frac{-\gamma}{2}\varphi^2 + \frac{1}{2}(\nabla\varphi)^2 + \frac{u\phi_0}{6}\varphi^3 + \frac{u}{4!}\varphi^4 \right]. \quad (10)$$

The constant term in H corresponds to the mean field result of its free-energy. After the shift the coefficient $-\gamma/2$ of the Gaussian terms in H is positive. Now the partition function and other physical quantities can be calculated in a similar way to that for the case $T > T_C$. The new φ^3 term in H shows the difference between the two phases in the transition. This term has no contribution to the partition function and two-point correlation function at first-order of the expansion, thus those quantities depend on $|\gamma|$ at first-order approximation. Due to the φ^3 term in H , the correlation functions among $2n + 1$ points, which are zero for high temperature phase, turn out to be nonzero up to the first-order approximation.

Above discussions allow a systematic study of the critical properties of finite-size systems with a one-component order parameter. Clearly, the techniques can be generalized to the case with n -component order parameter, which will be discussed elsewhere.

As a summary, a new perturbation theory of ϕ^4 model with a one-component order parameter is proposed to study finite-size effects of system near critical point. The new method provides a simple way for the calculation of the partition functions and correlation functions etc., and can be generalized to more complicated and realistic models.

This work was supported in part by the NNSF, the Hubei SF and the SECF in China.

REFERENCES

- [1] Erik Luijten and Henk W.J. Blöte, Phys. Rev. Lett. **76**, 1557 (1996); T. Ambrose and C.L. Chien, Phys. Rev. Lett. **76**, 1743 (1996); M. Krech and S. Dietrich, Phys. Rev. Lett. **66**, 345 (1991); Paul. A. Pearce and Andreas Klümper, Phys. Rev. Lett. **66**, 974 (1991).
- [2] M.E. Fisher, in *Critical Phenomena*, International School of Physics “ Enrico Fermi”, course 51, ed. M.S. Green’s (Academic, NewYork, 1971); M.E. Fisher and M.N. Baeber, Phys. Rev. Lett. **28**, 1516 (1972); K. Binder, Z. Phys. **B43**, 119 (1981).
- [3] M. Suzuki, Prog. Theor. Phys. **58**, 1142 (1977); E. Brézin, J. Physique **43**, 15 (1982).
- [4] E. Brézin and J. Zinn-Justin, Nucl. Phys. **B257**, 867 (1985).
- [5] J. Rudnick, H. Guo and D. Jasnow, J. Stat. Phys. **41**, 353 (1985).
- [6] E. Eisenriegler and R. Tomaschitz, Phys. Rev. **B35**, 4876 (1987); W. Huhn and V. Dohm, Phys. Rev. Lett. **61**, 1368 (1988); V. Dohm, Z. Phys. **B75**, 109 (1989); M. Krech and S. Dietrich, Phys. Rev. **A46**, 1886 (1992); 1922 (1992); P. Sutter and V. Dohm, Physica **B194-196**, 613 (1994).
- [7] R. Schmolke, A. Wacker, V. Dohm and D. Frank, Physica **B165& 166**, 575 (1990); J.J. Morris, J. Stat. Phys. **69**, 539 (1992).
- [8] R.C. Hwa and M.T. Nazirov, Phys. Rev. Lett. **69**, 741 (1992); R.C. Hwa and J. Pan, Phys. Lett. **B297**, 35 (1992); R.C. Hwa, Phys. Rev. **D47**, 2773 (1993); R.C. Hwa and J.Pan, Phys. Rev. **C50**, 383 (1994); A.K. Mohanty and S.K. Kataria, Phys. Rev. Lett. **73**, 2672 (1994); X. Cai, C.B. Yang and Z.M. Zhou, Phys. Rev. **C54**, No. 5 (1996).
- [9] A. Esser, V. Dohm and X.S. Chen, Physica **A222**, 355 (1995); X.S. Chen, V. Dohm and N. Schultka, Phys. Rev. Lett. **77**, 3641 (1996).